

CONNECTIONS ON NON-SYMMETRIC (GENERALIZED) RIEMANNIAN MANIFOLD AND GRAVITY

STEFAN IVANOV AND MILAN ZLATANOVIĆ

ABSTRACT. Connections with (skew-symmetric) torsion on non-symmetric Riemannian manifold satisfying the Einstein metricity condition (NGT with torsion) are considered. It is shown that an almost Hermitian manifold is an NGT with torsion if and only if it is a Nearly Kähler manifold. In the case of an almost contact metric manifold the NGT with torsion spaces are characterized and a possibly new class of almost contact metric manifolds is extracted. Similar considerations lead to a definition of a particular classes of almost para-Hermitian and almost paracontact metric manifolds. The conditions are given in terms of the corresponding Nijenhuis tensors and the exterior derivative of the skew-symmetric part of the non-symmetric Riemannian metric.

CONTENTS

1. Introduction	1
1.1. Motivation from general relativity	2
2. The geometric model	4
2.1. Linear connections on generalized Riemannian manifolds	4
2.2. Metric connections on generalized Riemannian manifold	6
2.3. The torsion tensor, skew-symmetric torsion	7
2.4. Eisenhart condition	8
2.5. Skew-symmetric torsion. Examples	8
3. Einstein metricity condition (NGT)	10
3.1. NGT involving the Nijenhuis tensor	11
3.2. NGT and skew-symmetric torsion	12
3.3. Almost Hermitian manifolds, $A^2 = -1$	12
3.4. Almost para-Hermitian manifolds, $A^2 = 1$	13
3.5. Almost contact metric structures.	14
3.6. Almost-nearly cosymplectic structures	16
3.7. Almost paracontact metric structures	16
References	17

1. INTRODUCTION

In this note we consider connections on a non-symmetric Riemannian manifold $(M, G = g + F)$ with a general assumption, coming from the non-symmetric gravitational theory, that the symmetric part g of G is non-degenerate. There are many examples of generalized Riemannian manifolds among which it deserves to mention the almost Hermitian, the almost contact, the almost para-Hermitian, the almost paracontact manifolds etc.

It is happened that the torsion and the covariant derivative of the non-symmetric metric determine a linear connection. The connection is unique and it is completely determined by the torsion and the Levi-Civita connection of the symmetric part of the non-symmetric metric. The Levi-Civita covariant derivative of the skew-symmetric part of the non-symmetric metric and the torsion supply a necessary condition for the existence of the connection.

1.1. Motivation from general relativity. General relativity (GR) was developed by A. Einstein in 1916 ^{Ein1} [9], with contributions by many others after 1916. In GR the equation $ds^2 = g_{ij}dx^i dx^j$, ($g_{ij} = g_{ji}$) is valid, where g_{ij} are functions of the point in the space. In GR which is the four dimensional space-time continuum metric properties depend on the mass distribution. The magnitudes g_{ij} are known as *gravitational potential*. Christoffel symbols which are commonly expressed by Γ_{ij}^k , play the role of the magnitudes which determine gravitational force field. General relativity explains gravity as the curvature of space-time.

In the GR the metric tensor is related by the Einstein equations $R_{ij} - \frac{1}{2}Rg_{ij} = T_{ij}$, where R_{ij} is the Ricci tensor of metric of space time, R is the scalar curvature of the metric, and T_{ij} is the energy-momentum tensor of matter. In 1922, A. Friedmann ^{Fried} [15] found a solution in which the universe may expand or contract, and later G. Lemaître ^{Lema} [32] derived a solution for an expanding universe. However, Einstein believed that the universe was apparently static, and since a static cosmology was not supported by the general relativistic field equations, he added a cosmological constant Λ to the field equations, which became $R_{ij} - \frac{1}{2}Rg_{ij} + \Lambda g_{ij} = T_{ij}$.

Since 1923 to the end of his life Einstein worked on various variants of Unified Field Theory (Non-symmetric Gravitational Theory-NGT) ^{Ein} [10]. This theory had the aim to unite the gravitation theory, to which is related GR, and the theory of electromagnetism. Introducing different variants of his NGT, Einstein used a complex basic tensor, with symmetric real part and skew-symmetric imaginary part. Beginning with 1950. Einstein used real non-symmetric basic tensor G , sometimes called *generalized Riemannian metric/manifold*).

Remark that at NGT the symmetric part g_{ij} of the basic tensor G_{ij} ($G_{ij} = g_{ij} + F_{ij}$) is related to gravitation, and skew-symmetric one F_{ij} to electromagnetism. The same is valid for the symmetric part of the connection and torsion tensor, respectively.

More recently the idea of a non-symmetric metric tensor appears in Moffat's non-symmetric gravitational theory ^{Mof} [35]. In Moffat's theory the skew-symmetric part of the metric tensor represents a Proca field (massive Maxwell field) which is a part of the gravitational interaction, contributing to the rotation of galaxies.

While on a Riemannian space the connection coefficients are expressed by virtue of the metric, g_{ij} , at Einstein's works on NGT the connection between these magnitudes is determined by the so called *Einstein metricity condition* i.e. the non-symmetric metric tensor G and the connection components Γ_{ij}^k are connected with the equations

$$\boxed{\text{metein}} \quad (1.1) \quad \frac{\partial G_{ij}}{\partial x^m} - \Gamma_{im}^p G_{pj} - \Gamma_{mj}^p G_{ip} = 0.$$

A generalized Riemannian manifold satisfying the Einstein metricity condition (^{metein} (1.1)) is also called NGT-space ^{Ein, Lof, Mof} [10, 34, 35].

The choice of a connection in NGT is not uniquely determined. In particular, in NGT there exist two kinds of covariant derivative, for example for tensor a_j^i :

$$a_{j|m}^+ = \frac{\partial a_j^i}{\partial x^m} + \Gamma_{pm}^i a_j^p - \Gamma_{jm}^p a_p^i; \quad a_{j|m}^- = \frac{\partial a_j^i}{\partial x^m} + \Gamma_{mp}^i a_j^p - \Gamma_{mj}^p a_p^i,$$

We investigate connections on a generalized Riemannian manifold $(M, G = g + F)$ with a general assumption, coming from NGT, that the symmetric part g of G is non-degenerate. There are many examples of generalized Riemannian manifolds such as the almost Hermitian, the almost contact, the almost para-Hermitian, the almost paracontact manifolds etc. In these cases the skew-symmetric part of G is played by the fundamental 2-form F .

We show that the torsion and the covariant derivative of G determine a unique linear connection ∇ with given torsion T and covariant derivative ∇G provided the covariant derivative of the skew-symmetric part F of G , ∇F satisfies certain compatibility condition which we expressed in terms of the torsion and the covariant derivative of the symmetric part g of G , ∇g (Theorem ^{gmain} 2.2). The connection is unique and it is completely determined from the symmetric part g of G .

We look for a linear connections with torsion preserving the non-symmetric tensor G . Surprisingly, we find that a linear connection ∇ preserves the non-symmetric metric, $\nabla G = 0$ if and only if it preserves its symmetric and skew-symmetric parts, $\nabla g = \nabla F = 0$. The last condition leads to a necessary constrain, i.e. a PDE for the skew-symmetric part F of G expressed in terms of the torsion. We show that there exists a unique connection preserving G provided its torsion is given and the mentioned above PDE is satisfied (c.f. Theorem ^{main} 2.3).

We paid a special attention when the torsion of the connection preserving the generalized Riemannian metric G is totally skew-symmetric with respect to the symmetric part g of G . One reason for that comes from the supersymmetric string theories and non-linear σ -models (see e.g. ^{Str, BBDG, BBE, BBDP, y4, BSethi, GKMW, GMPW, GMW, GKP, GPap} [41, 2, 3, 4, 5, 6, 17, 18, 19, 20, 21] and references therein) as well as from gravity theory itself ^{Ham} [26]. We find a condition in terms of the Nijenhuis tensor and the exterior derivative of the skew-symmetric part F of G and show that this condition is necessary and sufficient for the existence of connection with skew-symmetric torsion preserving the generalized Riemannian metric G (c.f. Theorem ^{skew} 2.4). We note that such a conditions were previously known on almost Hermitian, almost contact, almost para-Hermitian, almost paracontact manifolds ^{F1, GKMW, Zam} [14, 17, 43]. We derive the results there as a consequence of our considerations.

In Section ^{ngt} 3 we restrict our considerations to the NGT, i.e. a generalized Riemannian manifold satisfying the Einstein metricity condition (^{metein} 1.1). We present a condition in terms of the Nijenhuis tensor and the exterior derivative of F which guarantees the existence of a unique linear connection with skew-symmetric torsion preserving the NGT structure and show that the torsion is equal to minus one third of the exterior derivative dF of F (Theorem ^{pp1} 3.1).

One of the main contributions is the application of this result to the almost Hermitian and almost contact metric manifolds.

A careful analysis of our general condition for the existence of a connection with skew-symmetric torsion satisfying the Einstein metricity condition in the case of almost Hermitian manifold allow us to conclude that an almost Hermitian manifold satisfies the Einstein metricity condition with respect to a connection with skew-symmetric torsion if and only if it is a Nearly Kähler manifold (Theorem ^{nika} 3.2). In other words, an almost Hermitian manifold is NGT with skew-symmetric torsion exactly when it is a Nearly Kähler manifold. In this case the connection coincides with the Gray connection ^{Gr1, Gr2, Gr3} [22, 23, 24] which is the unique connection with skew-symmetric torsion preserving the Nearly Kähler structure ^{F1} [14]. The Nearly Kähler manifolds (called almost Tachibana spaces in ^{Yano} [42]) were developed by A. Gray ^{Gr1, Gr2, Gr3} [22, 23, 24] and intensively studied since then in ^{K1, N1, MNS, N2, But, FIMU} [30, 37, 36, 38, 8, 12]. Nearly Kähler manifolds appear also in supersymmetric string theories (see e.g. ^{P01, PS, LNP, GLNP} [39, 40, 31, 25]). The first complete inhomogeneous examples of 6-dimensional Nearly Kähler manifolds were presented recently in ^{Fosh} [13].

Applying the general condition for the existence of a connection with skew-symmetric torsion satisfying the Einstein metricity condition to the almost contact metric manifold we arrived at a

possibly new class of almost contact metric manifolds, which we call here *almost-nearly cosymplectic*. We show that this is the precise class of almost contact metric manifold which are NGT with skew-symmetric torsion, i.e. admitting a connection with skew-symmetric torsion satisfying the Einstein metricity condition. The simplest example of almost-nearly cosymplectic manifold is the trivial circle bundle over a (compact) nearly Kähler manifold.

The cases of almost para-Hermitian and almost paracontact metric manifold are also treated with respect to the NGT space with skew-symmetric torsion. We show that an almost para-Hermitian manifold admits an NGT connection with totally skew symmetric torsion if and only if it admits a linear connection preserving the almost para-Hermitian structure with a totally skew-symmetric torsion. We extract a possibly new class of almost paracontact metric structures which are NGT with skew-symmetric torsion. We characterized this class by an explicit formula for the covariant derivative with respect to the Levi-Civita connection of the fundamental two form.

2. THE GEOMETRIC MODEL

geomod

The fundamental (0,2) tensor G in non-symmetric (generalized) Riemannian manifold (M, G) is in general non-symmetric. It is decomposed in two parts, the symmetric part g and the skew-symmetric part F , $G(X, Y) = g(X, Y) + F(X, Y)$, where

metric

$$(2.1) \quad g(X, Y) = \frac{1}{2}(G(X, Y) + G(Y, X)), \quad F(X, Y) = \frac{1}{2}(G(X, Y) - G(Y, X)).$$

We assume that the symmetric part is non-degenerate of arbitrary signature. Therefore, we obtain a well defined (1,1) tensor A determined by the condition

m1

$$(2.2) \quad F(X, Y) = g(AX, Y).$$

We look for a natural linear connections ∇ preserving the generalized Riemannian metric G , $\nabla G = 0$ with torsion $T(X, Y) = \nabla_X Y - \nabla_Y X - [X, Y]$.

Convention 2.1. *In the whole paper we shall use the capital Latin letters $X, Y, \dots$ to denote smooth vector fields on a smooth manifold M which commute, $[X, Y] = 0$. Hence, $T(X, Y) = \nabla_X Y - \nabla_Y X$.*

The Levi-Civita connection corresponding to the symmetric non-degenerate (0,2) tensor g we denote with ∇^g . The Koszul formula reads

lcg

$$(2.3) \quad g(\nabla_X^g Y, Z) = \frac{1}{2} [Xg(Y, Z) + Yg(X, Z) - Zg(Y, X)].$$

We denote the (0,3) torsion tensor with respect to g by the same letter, $T(X, Y, Z) := g(T(X, Y), Z)$.

2.1. Linear connections on generalized Riemannian manifolds. In this section we show that a linear connection ∇ with torsion tensor T on a generalized Riemannian manifold is completely determined by the torsion and the covariant derivative ∇g of the symmetric part g of G . More precisely, we have

gmain

Theorem 2.2. *Let $(M, G = g + F)$ be a generalized Riemannian manifold and ∇^g be the Levi-Civita connection of g . Let ∇ be a linear connection with torsion T and denote the covariant derivative of the symmetric part g of G by ∇g . Then ∇ is unique determined by the following formula*

cgencon

$$(2.4) \quad g(\nabla_X Y, Z) = g(\nabla_X^g Y, Z) + \frac{1}{2} [T(X, Y, Z) + T(Z, X, Y) - T(Y, Z, X)] \\ - \frac{1}{2} [(\nabla_X g)(Y, Z) + (\nabla_Y g)(Z, X) - (\nabla_Z g)(Y, X)].$$

The covariant derivative ∇F of the skew-symmetric part F of G is given by

$$\boxed{\text{cconn2}} \quad (2.5) \quad (\nabla_X F)(Y, Z) = (\nabla_X^g F)(Y, Z) + \frac{1}{2} \left[T(X, Y, AZ) + T(Z, X, AY) \right] \\ + \frac{1}{2} \left[T(AZ, X, Y) + T(AZ, Y, X) + T(X, AY, Z) + T(Z, AY, X) \right] \\ + \frac{1}{2} \left[(\nabla_X g)(AY, Z) - (\nabla_X g)(Y, AZ) - (\nabla_Y g)(AZ, X) \right] \\ + \frac{1}{2} \left[(\nabla_Z g)(AY, X) + (\nabla_{AZ} g)(Y, X) - (\nabla_{AY} g)(Z, X) \right].$$

In particular, the exterior derivative dF of F satisfies

$$\text{cconn1} \quad (2.6) \quad dF(X, Y, Z) = -T(X, Y, AZ) - T(Y, Z, AX) - T(Z, X, AY) \\ + (\nabla_X F)(Y, Z) + (\nabla_Y F)(Z, X) + (\nabla_Z F)(X, Y).$$

(2.4) cgencon Conversely, any three tensors $T, \nabla g, \nabla F$ satisfying (2.5) lconn2 determine a unique linear connection via

Proof. It follows from (2.2) that the (1,1) tensor A is skew-symmetric with respect to the pseudo-Riemannian metric g , $g(AX, Y) = -g(X, AY)$. A simple calculation using (2.2) yields

$$\begin{aligned} \text{a2} \quad (2.7) \quad (\nabla_X F)(Y, Z) &= Xg(AY, Z) - g(A\nabla_X Y, Z) - g(AY, \nabla_X Z) \\ &= (\nabla_X g)(AY, Z) + g((\nabla_X A)Y, Z). \end{aligned}$$

From the definition of the covariant derivative of a (0,2) tensor, we have

$$\boxed{\text{covG}} \quad (2.8) \quad \begin{aligned} (\nabla_X G)(Y, Z) &= XG(Y, Z) - G(\nabla_X Y, Z) - G(Y, \nabla_X Z); \\ (\nabla_Y G)(Z, X) &= YG(Z, X) - G(\nabla_Y Z, X) - G(Z, \nabla_Y X); \\ (\nabla_Z G)(Y, X) &= ZG(Y, X) - G(\nabla_Z Y, X) - G(Y, \nabla_Z X). \end{aligned}$$

Sum the first two equalities and subtract the third equality to get using (2.1) and the definition of the torsion that

$$\boxed{\text{ccon1}} \quad (2.9) \quad (\nabla_X G)(Y, Z) + (\nabla_Y G)(Z, X) - (\nabla_Z G)(Y, X) = Xg(Y, Z) + Yg(Z, X) - Zg(Y, X) \\ + dF(X, Y, Z) - 2g(\nabla_X Y, Z) + G(Z, T(X, Y)) - G(Y, T(X, Z)) - G(T(Y, Z), X).$$

Using the Koszul formula (2.3), we obtain from (2.9) that

$$\boxed{\text{ccon2}} \quad (2.10) \quad 2g(\nabla_X Y, Z) = 2g(\nabla_X^g Y, Z) + dF(X, Y, Z) + G(Z, T(X, Y)) \\ - G(Y, T(X, Z)) - G(T(Y, Z), X) - (\nabla_X G)(Y, Z) - (\nabla_Y G)(Z, X) + (\nabla_Z G)(Y, X).$$

The skew-symmetric part with respect to X, Y of (2.10) gives precisely the equation (2.6).
 Substitute (2.6) into (2.10) and use (2.1) to get (2.4). Insert the already proved (2.4) into

$$(\nabla_X F)(Y, Z) = XF(Y, Z) - F(\nabla_X Y, Z) - F(Y, \nabla_X Z),$$

use [\(2.2\)](#) and [\(2.7\)](#) to obtain [\(2.5\)](#).

To complete the proof, observe that $\frac{\text{cconn2}}{(2.5)}$ implies $\frac{\text{cconn1}}{(2.6)}$. Indeed, taking the cyclic sum of $\frac{\text{cconn2}}{(2.5)}$ we get $\frac{\text{cconn1}}{(2.6)}$.

The converse follows from (2.4) by straightforward computation.

2.2. Metric connections on generalized Riemannian manifold. Here we investigate the existence of metric connections on a generalized Riemannian manifold. We have

main **Theorem 2.3.** *Let $(M, G = g + F)$ be a generalized Riemannian manifold and ∇^g be the Levi-Civita connection of g .*

- a) *A linear connection ∇ preserves the generalized Riemannian metric G if and only if it preserves its symmetric part g and its skew-symmetric part F , $\nabla G = 0 \Leftrightarrow \nabla g = \nabla F = 0 \Leftrightarrow \nabla A = 0$.*
- b) *If there exists a linear connection ∇ preserving the generalized Riemannian metric G , $\nabla G = 0$ with torsion T then the following condition holds*

conn2 (2.11)
$$(\nabla_X^g F)(Y, Z) = -\frac{1}{2} \left[T(X, Y, AZ) + T(Z, X, AY) \right] \\ - \frac{1}{2} \left[T(AZ, X, Y) + T(AZ, Y, X) + T(X, AY, Z) + T(Z, AY, X) \right].$$

In particular, the exterior derivative of F satisfies the next equality

conn1 (2.12)
$$dF(X, Y, Z) = F(T(X, Y), Z) + F(T(Y, Z), X) + F(T(Z, X), Y), \quad \text{equivalently} \\ dF(X, Y, Z) = -T(X, Y, AZ) - T(Y, Z, AX) - T(Z, X, AY);$$

*Conversely, if the condition **conn2** (2.11) is valid then there exists a unique linear connection ∇ with torsion T preserving the generalized Riemannian metric G determined by the torsion T with the formula*

gencon (2.13)
$$g(\nabla_X Y, Z) = g(\nabla_X^g Y, Z) + \frac{1}{2} \left[T(X, Y, Z) + T(Z, X, Y) - T(Y, Z, X) \right].$$

Proof. Suppose we have $0 = \nabla G = \nabla g + \nabla F$. The cyclic sum of this equality yields

$$(\nabla_X g)(Y, Z) + (\nabla_Y g)(Z, X) + (\nabla_Z g)(X, Y) \\ = -(\nabla_X F)(Y, Z) - (\nabla_Y F)(Z, X) - (\nabla_Z F)(X, Y).$$

Observe that the left hand side is totally symmetric while the right hand side is totally skew-symmetric which leads to the vanishing of each side,

new1 (2.14)
$$(\nabla_X g)(Y, Z) + (\nabla_Y g)(Z, X) + (\nabla_Z g)(X, Y) = 0; \\ (\nabla_X F)(Y, Z) + (\nabla_Y F)(Z, X) + (\nabla_Z F)(X, Y) = 0.$$

On the other hand, the symmetric part of ∇G with respect to the first two arguments gives

$$(\nabla_X g)(Y, Z) + (\nabla_Y g)(X, Z) + (\nabla_X F)(Y, Z) + (\nabla_Y F)(X, Z) = 0$$

which combined with the first identity in **new1** (2.14) yields

new2 (2.15)
$$(\nabla_Z g)(X, Y) = (\nabla_X F)(Y, Z) + (\nabla_Y F)(X, Z).$$

Substitute **new2** (2.15) into $\nabla G = \nabla g + \nabla F = 0$ to get

$$(\nabla_X F)(Y, Z) + (\nabla_Y F)(X, Z) + (\nabla_Z F)(X, Y) = 0$$

which combined with the second equality in **new1** (2.14) gives $2(\nabla_Y F)(X, Z) = 0$. Now $\nabla g = 0$ follows from **new2** (2.15). Clearly, the conditions $\nabla g = \nabla F = 0$ imply $\nabla G = 0$. We conclude from **a2** (2.7) that a metric connection ∇ , $\nabla g = 0$ preserves the two form F if and only if it preserves the (1,1) tensor A , $\nabla F = 0 \Leftrightarrow \nabla A = 0$ since g is non-degenerate. This proofs a).

Set $\nabla g = \nabla F = 0$ into Theorem **main** 2.2 to conclude b). Apply Theorem **main** 2.2 to complete the proof. $\square$

2.3. The torsion tensor, skew-symmetric torsion. It follows from Theorem ^{main}2.3 that any linear connection preserving the generalized Riemannian metric is completely determined by the torsion tensor T . In this section we investigate the torsion tensor. To this end we recall the definition of the Nijenhuis tensor N of the (1,1) tensor A (see e.g. ^{KN}[33]),

$$\text{nuj2} \quad (2.16) \quad N(X, Y) = [AX, AY] + A^2[X, Y] - A[AX, Y] - A[X, AY].$$

The Nijenhuis tensor is skew-symmetric by definition and it plays a fundamental role in almost complex (resp. almost para-complex) geometry. If $A^2 = -1$ (resp. $A^2 = 1$) then the celebrated Nulander-Nirenberg theorem (see, e.g. ^{KN}[33]) shows that an almost complex structure is integrable if and only if the Nijenhuis tensor vanishes.

Let ∇ be a linear connection preserving the generalized Riemannian metric $G, \nabla G = 0$. The ∇ preserves g, F and $A, \nabla g = \nabla F = \nabla A = 0$. Using the definition of the torsion and the covariant derivative ∇A we can express the Nijenhuis tensor in terms of the torsion and ∇A as follows

$$\text{nuj1} \quad (2.17) \quad N(X, Y) = (\nabla_{AX} A)Y - (\nabla_{AY} A)X - A(\nabla_X A)Y + A(\nabla_Y A)X \\ - T(AX, AY) - A^2 T(X, Y) + AT(AX, Y) + AT(X, AY).$$

We denote the Nijenhuis tensor of type (0,3) with respect to g with the same letter, $N(X, Y, Z) := g(N(X, Y), Z)$. Set $\nabla A = 0$ into ^{nuj1}(2.17) and use ⁿⁱ¹(2.2) to get

$$\text{nuj2} \quad (2.18) \quad N(X, Y, Z) = -T(AX, AY, Z) - T(X, Y, A^2 Z) - T(AX, Y, AZ) - T(X, AY, AZ).$$

2.3.1. The skew-symmetric torsion. Linear connections preserving a (pseudo) Riemannian metric and having totally skew-symmetric torsion i.e. the torsion satisfies the condition

$$\text{st} \quad (2.19) \quad T(X, Y, Z) = -T(X, Z, Y),$$

become very attractive in the last twenty years mainly due to the relations with supersymmetric string theories (see e.g. ^{Str,GMW,GPap,v4,Bsethi}[41, 19, 21, 5, 6] and references therein, for a mathematical treatment consult the nice overview ^{Apr}[1]). The main point is that the number of preserved supersymmetries is equal to the number of parallel spinors with respect to such a connection. This property reduces the holonomy group of the connection to be a subgroup of a group which is stabilizer of a non-trivial spinor such as $SU(n), Sp(n), G_2, Spin(7)$. It is happened that in that cases such a connection is unique ^{Str,GMW,FI,Iv}[41, 17, 14, 28] determined entirely by the structure induced by the parallel spinor.

We investigate when a generalized Riemannian manifold admits a metric connection with skew-symmetric torsion. We have

skew **Theorem 2.4.** *Let $(M, G = g + F)$ be a generalized Riemannian manifold and ∇^g be the Levi-Civita connection of g . If there exists a linear connection ∇ preserving the generalized Riemannian metric $G, \nabla G = 0$ with totally skew-symmetric torsion T then the following condition holds*

$$\text{ndf1} \quad (2.20) \quad N(X, Y, AZ) + N(X, Z, AY) = dF(X, Y, A^2 Z) + dF(X, Z, A^2 Y).$$

The torsion tensor satisfies the equality

$$\text{tor1} \quad (2.21) \quad T(AX, AY, Z) = -N(X, Y, Z) + dF(X, Y, AZ); \\ T(AX, Y, Z) = 2(\nabla_X^g F)(Y, Z) - dF(X, Y, Z).$$

The torsion connection ∇ is determined by the formula

$$\text{skct} \quad (2.22) \quad g(\nabla_X Y, Z) = g(\nabla_X^g Y, Z) + \frac{1}{2}T(X, Y, Z).$$

If in addition the skew-symmetric part F of the generalized Riemannian metric G is closed, $dF = 0$, then ^{ndf1}(2.20) and ^{tor1}(2.21) hold inserting $dF = 0$.

Proof The equalities (2.21) follow by a straightforward calculations from (2.19), (2.18), (2.12) and (2.11). Now, the equality (2.20) is an easy consequence from the first equality in (2.21). The formula (2.22) follows from (2.13) and (2.19). $\square$

2.4. Eisenhart condition. Eisenhart [Eis] was one of the first who proposed a connection with skew-symmetric torsion to be applied in general relativity. In the sense of Eisenhart's definition (see [Eis]) generalized Riemannian manifold is a differentiable manifold with a non-symmetric basic tensor $G(X, Y) = g(X, Y) + F(X, Y)$ with a connection explicitly defined by the equation

$$\text{Eisen1} \quad (2.23) \quad g(\nabla_X Y, Z) = \frac{1}{2} [XG(Y, Z) + YG(Z, X) - ZG(Y, X)].$$

It is easy to see that (Eisen1 (2.23)) can be written in the form

$$g(\nabla_X Y, Z) = \frac{1}{2} [XG(Y, Z) + YG(Z, X) - ZG(Y, X)] = g(\nabla_X^g Y, Z) + \frac{1}{2} dF(X, Y, Z)$$

which shows that the symmetric part g of G is covariantly constant, $\nabla g = 0$ and the torsion T is totally skew-symmetric determined by the equation $T(X, Y, Z) = dF(X, Y, Z)$. Using relations (2.7) and (2.17), we calculate that the Nijenhuis tensor satisfies

$$\text{EisenhN1} \quad (2.24) \quad N(X, Y, Z) = (\nabla_{AX}^g F)(Y, Z) - (\nabla_{AY}^g F)(X, Z) + (\nabla_X^g F)(Y, AZ) - (\nabla_Y^g F)(X, AZ).$$

If, in addition, the Eisenhart connection preserves the generalized Riemannian metric G , $\nabla g = \nabla F = 0$, then equation (EisenhN1 (2.24)) reduces to (2.18) with $T = dF$.

2.5. Skew-symmetric torsion. Examples. A generalized Riemannian metric G is equivalent to a choice of a pseudo-Riemannian metric g and a 2-form F (an (1,1) tensor A satisfying (2.2)), such that $G = g + F$. A generalized metric connection, i.e. a linear connection preserving G , is a Riemannian connection preserving the 2-form F or equivalently, a Riemannian connection preserving the (1,1)-tensor A . This supplies a number of examples.

2.5.1. Almost Hermitian manifolds, $A^2 = -1$. Let us consider an almost Hermitian manifold (M, g, A) , i.e. Riemannian manifold (M, g) of dimension $n (= 2m \geq 4)$ endowed with an almost complex structure, the endomorphism A satisfies $A^2 = -I$, $F(X, Y) = g(AX, Y)$, $g(AX, AY) = g(X, Y)$. The 2-form F is the Kähler form (note the sign difference of the Kähler form in [14]).

In this case the Nijenhuis tensor (2.16) has the properties

$$\text{propAHN1} \quad (2.25) \quad N(AX, Y, Z) = N(X, AY, Z) = N(X, Y, AZ)$$

Then Theorem 2.4 gives the following well known result

Corollary 2.5 ([41, 16, 14]). *On an almost hermitian manifold (M, g, A) there exists a unique linear connection ∇ preserving the generalized Riemannian metric $G = g + F$ with totally skew-symmetric torsion T if and only if the Nijenhuis tensor is totally skew-symmetric, $N(X, Y, Z) = -N(X, Z, Y)$. The torsion T is determined by $T(X, Y, Z) = N(X, Y, Z) + dF(AX, AY, AZ)$. In particular, an almost Kähler manifold, $dF = 0$, admits such a connection if and only if it is a Kähler manifold, $N = 0$*

2.5.2. Almost para-hermitian manifolds, $A^2 = 1$. Almost para-Hermitian manifold (M, g, A) is Riemannian manifold (M, g) endowed with endomorphism A satisfying $A^2 = I$, $F(X, Y) = g(AX, Y)$, $g(AX, AY) = -g(X, Y)$. Such a manifold is of even dimension $2n$, the eigen-subbundles of the paracomplex structure A are of equal dimension n and the metric g is of neutral signature (n, n) . In this case the Nijenhuis tensor (2.16) has the properties (propAHN1 (2.25)) and Theorem 2.4 gives the following well known result (note the sign difference of the 2-form F in [29])

APHskew

Corollary 2.6 ([29]). *On an almost para-hermitian manifold (M, g, A) there exists a unique linear connection ∇ preserving the generalized Riemannian metric $G = g + F$ with totally skew-symmetric torsion T if and only if the Nijenhuis tensor is totally skew-symmetric, $N(X, Y, Z) = -N(X, Z, Y)$. The torsion T is determined by $T(X, Y, Z) = -N(X, Y, Z) + dF(AX, AY, AZ)$. In particular, an almost para-Kähler manifold, $dF = 0$, admits such a connection if and only if it is a para-Kähler manifold, $N = 0$*

2.5.3. Almost contact metric structures. We consider an almost contact metric manifold $(M^{2n+1}, g, A, \eta, \xi)$, i.e., a $(2n + 1)$ -dimensional Riemannian manifold equipped with a 1-form η , a $(1,1)$ -tensor A and a vector field ξ dual to η with respect to the metric g , $\eta(\xi) = 1, \eta(X) = g(X, \xi)$ such that the following compatibility conditions are satisfied (see e.g. [7])

$$(2.26) \quad A^2 = -id + \eta \otimes \xi, \quad g(AX, AY) = g(X, Y) - \eta(X)\eta(Y), \quad F(X, Y) = g(AX, Y), \quad A\xi = 0.$$

In this case the skew-symmetric part F of $G = g + F$ is degenerate, $F(\xi, X) = 0$ and has rank $2n$. Then Theorem 2.3 together with (2.26) implies $0 = g((\nabla_X A)AY, \xi) = (\nabla_X \eta)Y = g(\nabla_X \xi, Y)$, i.e. $\nabla \eta = \nabla \xi = 0$. Now (2.22) yields $0 = g(\nabla_X^g \xi, Y) + \frac{1}{2}T(X, \xi, Y)$ which shows that ξ is a Killing vector field because T is skew-symmetric and

aconeta

$$(2.27) \quad d\eta = \xi \lrcorner T, \quad \xi \lrcorner d\eta = 0,$$

where $\lrcorner$ is the interior multiplication.

We obtain from (2.18) applying (2.26) and (2.27) that

nujac

$$(2.28) \quad N(X, Y, Z) + \eta(Z)d\eta(X, Y) = -T(AX, AY, Z) + T(X, Y, Z) - T(AX, Y, AZ) - T(X, AY, AZ)$$

The right hand side of equation (2.28) is totally skew-symmetric which yields that the tensor

nujac0

$$(2.29) \quad N^{ac} = N + d\eta \otimes \eta$$

is totally skew-symmetric, $N^{ac}(X, Y, Z) = -N^{ac}(X, Z, Y)$. In fact, the tensor N^{ac} is usually called the Nijenhuis tensor in almost contact geometry (see e.g. [7]).

We get from (2.28) applying (2.26) and (2.27) that

nujac1

$$(2.30) \quad N(\xi, Y, Z) = d\eta(Y, Z) - d\eta(AY, AZ).$$

We calculate from (2.16) taking into account (2.26) and (2.27) that

acs

$$(2.31) \quad N(AX, AY, Z) = -N(X, Y, Z) - \eta(Z) \left[d\eta(AX, AY) + d\eta(X, Y) \right] + \eta(X)N(\xi, Y, Z) - \eta(Y)N(\xi, X, Z)$$

The first formula in (2.21) together with (2.27) and (2.31) yields

$$\begin{aligned} T(X, Y, Z) &= -N(AX, AY, Z) + dF(AX, AY, AZ) + \eta(X)d\eta(Y, Z) + \eta(Y)d\eta(Z, X) \\ &= N(X, Y, Z) + dF(AX, AY, AZ) + \eta(Z) \left[d\eta(AX, AY) + d\eta(X, Y) \right] - \eta(X)N(\xi, Y, Z) + \eta(Y)N(\xi, X, Z) \\ &\quad + \eta(X)d\eta(Y, Z) + \eta(Y)d\eta(Z, X). \end{aligned}$$

We obtain from this equality applying (2.30) the following formula for the skew-symmetric torsion

tac

$$(2.32) \quad T(X, Y, Z) = N(X, Y, Z) + \eta(Z)d\eta(X, Y) + dF(AX, AY, AZ) + \eta(Z)d\eta(AX, AY) + \eta(Y)d\eta(AZ, AX) + \eta(X)d\eta(AY, AZ).$$

Observe using $(\text{nuj})_{(2.16)}$, $(\text{acon})_{(2.26)}$ and $(\text{nujac0})_{(2.29)}$ that $d\eta(AX, AY) = -N(X, Y, \xi) = -N^{ac}(X, Y, \xi) + d\eta(X, Y)$. Substitute the last equality into $(\text{tac})_{(2.32)}$ and use the skew-symmetry of N^{ac} to get the equality obtained in $(\text{FI})_{[14]}$

$$(\text{tac1}) \quad (2.33) \quad T(X, Y, Z) = \eta \wedge d\eta + N^{ac} - d^A F - \eta(Z) \wedge (\xi \lrcorner N^{ac}), \quad d^A F(X, Y, Z) = -dF(AX, AY, AZ).$$

Theorem $(\text{skew})_{2.4}$ gives the next well known result (note the sign difference of the 2-form F in $(\text{FI})_{[14]}$)

AcHskew

Corollary 2.7 $(\text{FI})_{[14]}$. *On an almost contact metric manifold (M, g, A, F, η, ξ) there exists a unique linear connection ∇ preserving the generalized Riemannian metric $G = g + F$ with totally skew-symmetric torsion T if and only if the almost contact Nijenhuis tensor N^{ac} is totally skew-symmetric, $N^{ac}(X, Y, Z) = -N^{ac}(X, Z, Y)$ and ξ is a Killing vector field. The torsion T is determined by $(\text{tac1})_{(2.33)}$ or, equivalently by $(\text{tac})_{(2.32)}$.*

2.5.4. Almost paracontact metric structures. We consider an almost paracontact metric manifold $(M^{2n+1}, g, A, \eta, \xi)$, i.e., a $(2n+1)$ -dimensional pseudo Riemannian manifold of signature $(n+1, n)$ equipped with a 1-form η , a $(1,1)$ -tensor A and a vector field ξ dual to η with respect to the metric g , $\eta(\xi) = 1$, $\eta(X) = g(X, \xi)$ such that the following compatibility conditions hold (see e.g. $(\text{Zam})_{[43]}$)

$$(\text{apcon}) \quad (2.34) \quad A^2 = id - \eta \otimes \xi, \quad g(AX, AY) = -g(X, Y) + \eta(X)\eta(Y), \quad F(X, Y) = g(AX, Y), \quad A\xi = 0.$$

In this case the skew-symmetric part F of $G = g + F$ is degenerate, $F(\xi, X) = 0$ and has rank $2n$. Theorem $(\text{main})_{2.3}$ together with $(\text{apcon})_{(2.34)}$ implies $0 = g((\nabla_X A)AY, \xi) = -(\nabla_X \eta)Y = -g(\nabla_X \xi, Y)$, i.e. $\nabla \eta = \nabla \xi = 0$. Now $(\text{skct})_{(2.22)}$ yields $0 = g(\nabla_X^g \xi, Y) + \frac{1}{2}T(X, \xi, Y)$ which shows that ξ is a Killing vector field because T is skew-symmetric and $d\eta = \xi \lrcorner T$, $\xi \lrcorner d\eta = 0$. We obtain from $(\text{nuj2})_{(2.18)}$ applying $(\text{apcon})_{(2.34)}$ and the last equality

$$(\text{nujapc}) \quad (2.35) \quad N(X, Y, Z) - \eta(Z)d\eta(X, Y) = -T(AX, AY, Z) - T(X, Y, Z) - T(AX, Y, AZ) - T(X, AY, AZ)$$

The right hand side of equation $(\text{nujapc})_{(2.35)}$ is totally skew-symmetric which yields that the tensor $N^{apc} = N - d\eta \otimes \eta$ is totally skew-symmetric, $N^{apc}(X, Y, Z) = -N^{apc}(X, Z, Y)$. In fact, the tensor N^{apc} is usually called *the Nijenhuis tensor in almost paracontact geometry* (see e.g. $(\text{Zam})_{[43]}$).

As in the previous case, Theorem $(\text{skew})_{2.4}$ gives the following well known result (note the sign difference of the 2-form F in $(\text{Zam})_{[43]}$)

AcHskewp

Corollary 2.8 $(\text{Zam})_{[43]}$. *On an almost paracontact metric manifold (M, g, A, F, η, ξ) there exists a unique linear connection ∇ preserving the generalized Riemannian metric $G = g + F$ with totally skew-symmetric torsion T if and only if the almost paracontact Nijenhuis tensor N^{apc} is totally skew-symmetric, $N^{apc}(X, Y, Z) = -N^{apc}(X, Z, Y)$ and ξ is a Killing vector field. The torsion T is determined by*

$$T(X, Y, Z) = \eta \wedge d\eta - N^{apc} - d^A F + \eta(Z) \wedge (\xi \lrcorner N^{ac}), \quad d^A F(X, Y, Z) = -dF(AX, AY, AZ).$$

3. EINSTEIN METRICITY CONDITION (NGT)

ngt

In his attempt to construct an unified field theory (Non-symmetric Gravitational Theory, briefly NGT) A.Einstein $(\text{Ein})_{[10]}$ considered a generalized Riemannian manifold and use the so called metricity condition $(\text{metein})_{(1.1)}$ which can be written as follows $XG(Y, Z) - G(\nabla_Y X, Z) - G(Y, \nabla_X Z) = 0$, see also $(\text{Mof, LMof})_{[35, 34]}$ for a little bit more general right hand side of $(\text{metein})_{(1.1)}$.

In view of $(\text{covG})_{(2.8)}$, the definition of the torsion, $(\text{metric})_{(2.1)}$ and $(\text{m1})_{(2.2)}$ the metricity condition $(\text{metein})_{(1.1)}$ can be written in the form

metein1

$$(\text{metein1}) \quad (3.1) \quad (\nabla_X G)(Y, Z) = -G(T(X, Y), Z) \Leftrightarrow (\nabla_X (g + F))(Y, Z) = -T(X, Y, Z) + T(X, Y, AZ).$$

A general solution for the connection ∇ satisfying $(\text{metein } 1.1)$ is given in terms of g, F, T $(\text{Hlav } 2.7)$ (see also $(\text{Mor } 3.5)$). Here we show that a solution can be expressed in terms of the exterior derivative dF of F and find formulas for the covariant derivatives of ∇g and ∇F .

Taking the cyclic sum in $(\text{metein1 } 3.1)$ and applying $(\text{cconn1 } 2.6)$, we obtain

$$\begin{aligned} \text{ein2} \quad (3.2) \quad & (\nabla_X g)(Y, Z) + (\nabla_Y g)(Z, X) + (\nabla_Z g)(X, Y) \\ & = -dF(X, Y, Z) - T(X, Y, Z) - T(Y, Z, X) - T(Z, X, Y). \end{aligned}$$

It is easy to observe that the left hand side of $(\text{lein2 } 3.2)$ is symmetric while the right hand side is skew-symmetric. Hence, we get

$$\begin{aligned} \text{ein3} \quad (3.3) \quad & (\nabla_X g)(Y, Z) + (\nabla_Y g)(Z, X) + (\nabla_Z g)(X, Y) = 0; \\ & dF(X, Y, Z) = -T(X, Y, Z) - T(Y, Z, X) - T(Z, X, Y). \end{aligned}$$

The symmetric part of $(\text{metein1 } 3.1)$ with respect to X, Y gives

$$(\nabla_X g)(Y, Z) + (\nabla_Y g)(X, Z) + (\nabla_X F)(Y, Z) + (\nabla_Y F)(X, Z) = 0,$$

which combined with the first equality in $(\text{lein3 } 3.3)$ yields

$$\text{ein4} \quad (3.4) \quad (\nabla_Z g)(X, Y) = (\nabla_X F)(Y, Z) + (\nabla_Y F)(X, Z).$$

Substitute $(\text{lein4 } 3.4)$ into $(\text{metein1 } 3.1)$ and use $(\text{cconn1 } 2.6)$ to get

$$\text{ein5} \quad (3.5) \quad (\nabla_Z F)(X, Y) = \frac{1}{2} [dF(X, Y, Z) + T(X, Y, Z) - T(Z, Y, AX) + T(Z, X, AY)].$$

We obtain inserting $(\text{lein5 } 3.5)$ into $(\text{lein4 } 3.4)$ that

$$\text{ein6} \quad (3.6) \quad (\nabla_X g)(Y, Z) = -\frac{1}{2} [T(X, Y, Z) - T(X, Y, AZ) + T(X, Z, Y) - T(X, Z, AY)].$$

Applying $(\text{lein6 } 3.6)$ and the second equation in $(\text{lein3 } 3.3)$ we obtain from $(\text{cgencon } 2.4)$ that

$$\begin{aligned} \text{genconein} \quad (3.7) \quad & g(\nabla_X Y, Z) = g(\nabla_X^g Y, Z) + \frac{1}{2} [T(X, Y, Z) - T(X, Z, AY) - T(Y, Z, AX)] \\ & = g(\nabla_X^g Y, Z) - \frac{1}{2} [dF(X, Y, Z) + T(Z, X, Y) + T(Y, Z, X)] + \frac{1}{2} [T(Z, X, AY) + T(Z, Y, AX)]. \end{aligned}$$

3.1. NGT involving the Nijenhuis tensor. We are going to involve the Nijenhuis tensor in our attempt to determine the torsion T . We start with an expression of the covariant derivative ∇A . Substitute $(\text{lein5 } 3.5)$ and $(\text{lein6 } 3.6)$ into $(\text{a2 } 2.7)$, we find

$$\begin{aligned} \text{ein7} \quad (3.8) \quad & g((\nabla_X A)Y, Z) = (\nabla_X F)(Y, Z) - (\nabla_X g)(AY, Z) \\ & = \frac{1}{2} [dF(X, Y, Z) + T(Y, Z, X) + T(X, Y, AZ) + T(X, AY, Z) - T(X, AY, AZ) - T(X, Z, A^2 Y)] \end{aligned}$$

Substitute $(\text{lein7 } 3.8)$ into $(\text{hui1 } 2.17)$ to get after some calculations the following equality

$$\begin{aligned} \text{ein8} \quad (3.9) \quad & N(X, Y, Z) = dF(X, Y, AZ) + \frac{1}{2} [dF(AX, Y, Z) + dF(X, AY, Z)] \\ & + \frac{1}{2} [T(Y, Z, AX) - T(X, Z, AY) + T(Y, AZ, X) - T(X, AZ, Y) - T(X, AY, A^2 Z) - T(AX, Y, A^2 Z)] \\ & - \frac{1}{2} [T(Z, AY, A^2 X) - T(Z, AX, A^2 Y) + T(AZ, Y, A^2 X) - T(AZ, X, A^2 Y)] - T(AX, AY, AZ). \end{aligned}$$

3.2. NGT and skew-symmetric torsion. Now we consider the case of totally skew-symmetric torsion, $T(X, Y, Z) = -T(X, Z, Y)$. From the considerations above, we have

pp1 **Theorem 3.1.** *A generalized Riemannian manifold $(M, G = g + F)$ admits a linear connection satisfying the Einstein metricity condition (1.1) with totally skew-symmetric torsion T if and only if the Nijenhuis tensor N and the exterior derivative of F satisfy the relation*

skew1 (3.10)
$$N(X, Y, Z) = \frac{2}{3}dF(X, Y, AZ) + \frac{1}{3}dF(AX, Y, Z) + \frac{1}{3}dF(X, AY, Z) + \frac{1}{3}dF(AX, AY, AZ) \\ - \frac{1}{6}\left[dF(A^2X, Y, AZ) + dF(A^2X, AY, Z) + dF(X, A^2Y, AZ) - dF(X, AY, A^2Z)\right] \\ - \frac{1}{6}\left[dF(AX, A^2Y, Z) - dF(AX, Y, A^2Z)\right]$$

In this case the skew-symmetric torsion is completely determined by the exterior derivative of the skew-symmetric part of the generalized Riemannian metric,

tordfnew (3.11)
$$T(X, Y, Z) = -\frac{1}{3}dF(X, Y, Z),$$

the Einstein metricity condition has the form $(\nabla_X G)(Y, Z) = \frac{1}{3}\left[dF(X, Y, Z) - dF(X, Y, AZ)\right]$ and it is equivalent to the following two conditions

skew0 (3.12)
$$(\nabla_X g)(Y, Z) = -\frac{1}{6}\left[dF(X, Y, AZ) - dF(X, AY, Z)\right]; \\ (\nabla_X F)(Y, Z) = \frac{1}{6}\left[2dF(X, Y, Z) - dF(X, Y, AZ) - dF(X, AY, Z)\right].$$

The connection is unique determined by the formula

newnbl (3.13)
$$g(\nabla_X Y, Z) = g(\nabla_X^g Y, Z) - \frac{1}{6}dF(X, Y, Z) - \frac{1}{6}dF(X, AY, Z) + \frac{1}{6}dF(AX, Y, Z).$$

The covariant derivative of F and A with respect to the Levi-Civita connection ∇^g are given by

ff2 (3.14)
$$(\nabla_X^g F)(Y, Z) = g((\nabla_X^g A)Y, Z) \\ = \frac{1}{3}dF(X, Y, Z) + \frac{1}{3}dF(X, AY, AZ) - \frac{1}{6}dF(AX, Y, AZ) - \frac{1}{6}dF(AX, AY, Z).$$

Proof. The second equation of (3.3) together with the skew-symmetric property of the torsion imply (3.11). Using the already obtained (3.11) reduces (3.9) to (3.10). Now, (3.12) and (3.13) follow from (3.5), (3.6), (3.7) and (3.11). Applying (3.11) and (3.12) to (2.5), we obtain

ff1 (3.15)
$$(\nabla_X F)(Y, Z) = (\nabla_X^g F)(Y, Z) - \frac{1}{6}\left[dF(X, Y, AZ) + dF(Z, X, AY)\right] \\ - \frac{1}{6}\left[2dF(X, AY, AZ) + dF(Z, AY, AX) + dF(Y, AX, AZ)\right].$$

A substitution of the second formula in (3.12) into (3.15) gives (3.14).

For the converse, it is straightforward to check that the connection determining by (3.13) satisfies the required properties provided (3.10) holds. This completes the proof. $\square$

3.3. Almost Hermitian manifolds, $A^2 = -1$. The condition (3.10) in the almost hermitian case takes the form

skewah (3.16)
$$N(X, Y, Z) = dF(X, Y, AZ) + \frac{1}{3}dF(AX, Y, Z) + \frac{1}{3}dF(X, AY, Z) + \frac{1}{3}dF(AX, AY, AZ).$$

Using the properties of the Nijenhuis tensor of an almost Hermitian manifold, (^{propAHN1} (2.25)), we get from (^{skewah} (3.16)) that the exterior derivative dF of the Kähler form satisfies

$$\boxed{\text{dfah}} \quad (3.17) \quad dF(X, Y, AZ) = dF(X, AY, Z) = dF(AX, Y, Z) = -dF(AX, AY, AZ),$$

i.e. dF is of type $(3, 0) + (0, 3)$ with respect to the almost complex structure A . Substitute (^{dfah} (3.17)) into (^{skewah} (3.16)) to get $N(X, Y, Z) = \frac{4}{3}dF(X, Y, AZ)$.

At this point we recall that an almost Hermitian manifold is said to be Nearly Kähler if the covariant derivative of the almost complex structure A with respect to the Levi-Civita connection ∇^g of the metric g is skew-symmetric,

$$\boxed{\text{nk2}} \quad (3.18) \quad (\nabla_X^g A)X = 0 \quad \Leftrightarrow \quad (\nabla_X^g F)(X, Y) = 0.$$

The Nearly Kähler manifolds (called almost Tachibana spaces in ^{Yano} [42]) were developed by A. Gray ^{Gr1, Gr2, Gr3} [22, 23, 24] and intensively studied since then in ^{K1, N1, MNS, N2, But, FIMU} [30, 37, 36, 38, 8, 12]. Nearly Kähler manifolds in dimension 6 are Einstein manifolds with positive scalar curvature, the Nijenhuis tensor N is a 3-form and it is parallel with respect to the Gray characteristic connection (see ^{K1} [30]). This connection was defined by Gray ^{Gr1, Gr2, Gr3} [22, 23, 24] and it turns out to be the unique linear connection preserving the nearly Kähler structure and having totally skew-symmetric torsion (see ^{F1} [14]). Nearly Kähler manifolds appear also in supersymmetric string theories (see e.g. ^{Po1, PS, LNP, GLNP} [39, 40, 31, 25] etc).

We obtain from Theorem ^{pp1} 3.1 that

Theorem 3.2. *Let (M, A, g, F) be an almost Hermitian manifold with a Kähler 2-form F considered as a generalized Riemannian manifold $(M, G = g + F)$. Then (M, G) satisfies the Einstein metricity condition (^{metein} (1.1)) with a totally skew-symmetric torsion T if and only if it is a Nearly Kähler manifold.*

The skew-symmetric torsion is determined by the condition

$$\boxed{\text{nk3}} \quad (3.19) \quad T(X, Y, Z) = -\frac{1}{3}dF(X, Y, Z) = \frac{1}{4}N(X, Y, AZ).$$

The connection is unique given by the formula

$$g(\nabla_X Y, Z) = g(\nabla_X^g Y, Z) - \frac{1}{6}dF(X, Y, Z) = g(\nabla_X^g Y, Z) + \frac{1}{8}N(X, Y, AZ).$$

The Einstein metricity condition has the form $(\nabla_X G)(Y, Z) = \frac{1}{3}[dF(X, Y, Z) - dF(X, Y, AZ)]$. The covariant derivative of g and the Kähler form F are

$$(\nabla_X g)(Y, Z) = 0; \quad (\nabla_X F)(Y, Z) = \frac{1}{3}[dF(X, Y, Z) - dF(X, Y, AZ)].$$

Proof. Applying (^{nk3} (3.19)) and (^{dfah} (3.17)) to (^{cconn2} (2.5)) we conclude that the Nearly Kähler condition (^{nk2} (3.18)) holds. The rest of the theorem follow from Theorem ^{pp1} 3.1. $\square$

3.4. Almost para-Hermitian manifolds, $A^2 = 1$. The condition (^{skew1} (3.10)) in the almost para-hermitian case takes the form

$$N(X, Y, Z) = \frac{1}{3}dF(X, Y, AZ) + \frac{1}{3}dF(AX, Y, Z) + \frac{1}{3}dF(X, AY, Z) + \frac{1}{3}dF(AX, AY, AZ),$$

which implies that the Nijenhuis tensor is totally skew symmetric.

We obtain from Theorem ^{pp1} 3.1 and ^{F2} [29, Proposition 3.1]

Theorem 3.3. *Let (M, A, g, F) be an almost para-Hermitian manifold with a Kähler 2-form F considered as a generalized Riemannian manifold $(M, G = g + F)$. Then (M, G) satisfies the Einstein metricity condition (^{metein} (1.1)) with a totally skew-symmetric torsion T if and only if the Nijenhuis tensor is totally skew-symmetric.*

ngtskewp

The skew-symmetric torsion is determined by $T(X, Y, Z) = -\frac{1}{3}dF(X, Y, Z)$, the Einstein metric-ity condition has the form $(\nabla_X G)(Y, Z) = \frac{1}{3}[dF(X, Y, Z) - dF(X, Y, AZ)]$ and the equalities [\(3.12\)](#), [\(3.13\)](#) and [\(3.14\)](#) hold with $A^2 = 1$.

Remark 3.4. In particular, the nearly paraKähler manifolds defined by $(\nabla_X^g A)X = 0$ have totally skew symmetric Nijenhuis tensor, dF share the properties [\(2.25\)](#) of the Nijenhuis tensor and $N(X, Y, Z) = \frac{4}{3}dF(X, Y, AZ)$ [\[29, Proposition 5.1\]](#). In this case the NGT connection with skew symmetric torsion preserves the symmetric part of G , $\nabla g = 0$.

Remark 3.5. Combining Theorem [3.3](#) and Corollary [2.6](#) we conclude that an almost para-Hermitian manifold admits an NGT connection with totally skew symmetric torsion if and only if it admits a linear connection preserving the almost para-Hermitian structure with a totally skew-symmetric torsion.

3.5. Almost contact metric structures. Let us consider the case of an almost contact metric structure which is defined in the Section 4.3. The fact $g(\xi, \xi) = 1$ and [\(3.14\)](#) imply

$$\text{kill} \quad (3.20) \quad (\nabla_X^g \eta)Z = g(\nabla_X^g \xi, Z) = -g((\nabla_X^g A)\xi, AZ) = \frac{1}{3}dF(X, AZ, \xi) + \frac{1}{6}dF(AX, Z, \xi).$$

Consequently, we obtain

$$\text{xi1} \quad (3.21) \quad d\eta(X, Z) = \frac{1}{2}dF(X, AZ, \xi) + \frac{1}{2}dF(AX, Z, \xi), \quad d\eta(X, \xi) = 0, \quad d\eta(AX, Z) = d\eta(X, AZ).$$

Applying [\(2.26\)](#) and [\(3.21\)](#), we simplify [\(3.10\)](#) to get

$$\text{skewac} \quad (3.22) \quad N(X, Y, Z) = dF(X, Y, AZ) + \frac{1}{3}dF(AX, Y, Z) + \frac{1}{3}dF(X, AY, Z) + \frac{1}{3}dF(AX, AY, AZ) - \frac{1}{3}[d\eta(Y, Z)\eta(X) + d\eta(Z, X)\eta(Y) - d\eta(X, Y)\eta(Z)].$$

On the other hand, [\[7, Blair Lemma 6.1\]](#), in our notation, reads

$$\text{Problem} \quad (3.23) \quad 2g((\nabla_X^g A)Y, Z) = dF(X, Y, Z) - dF(X, AY, AZ) + N^{ac}(Y, Z, AX) + [d\eta(AY, Z) - d\eta(AZ, Y)]\eta(X) - d\eta(X, AY)\eta(Z) + d\eta(X, AZ)\eta(Y).$$

Substitute [\(3.14\)](#) into [\(3.23\)](#), use [\(3.21\)](#) and observe $N^{ac}(Y, Z, AX) = N(Y, Z, AX)$ to get

$$\text{skewacB} \quad (3.24) \quad N(Y, Z, AX) = -2d\eta(AY, Z)\eta(X) + d\eta(X, AY)\eta(Z) - d\eta(X, AZ)\eta(Y) - \frac{1}{3}dF(X, Y, Z) + \frac{5}{3}dF(X, AY, AZ) - \frac{1}{3}dF(AX, Y, AZ) - \frac{1}{3}dF(AX, AY, Z).$$

Setting $X = \xi$ into [\(3.24\)](#) gives $0 = -\frac{1}{3}dF(\xi, Y, Z) + \frac{5}{3}dF(\xi, AY, AZ) - 2d\eta(AY, Z)$ which, in view of [\(3.21\)](#), yields

$$\text{dfdet} \quad (3.25) \quad dF(Y, AZ, \xi) = d\eta(Y, Z) = dF(AY, Z, \xi).$$

Applying [\(3.25\)](#) to [\(3.20\)](#) leads to

$$\text{kill1} \quad (3.26) \quad (\nabla_X^g \eta)Y = g(\nabla_X^g \xi, Y) = \frac{1}{2}d\eta(X, Y),$$

which shows that the vector field ξ is a Killing vector field.

The equalities [\(3.22\)](#) and [\(3.25\)](#) imply

$$\text{nijxi} \quad (3.27) \quad N(X, Y, \xi) = N(\xi, X, Y) = d\eta(X, Y).$$

Applying [\(5.27\)](#), we obtain from [\(5.24\)](#)

$$\begin{aligned} (3.28) \quad N(X, Y, Z) &= d\eta(Y, Z)\eta(X) + \frac{5}{3}d\eta(X, Y)\eta(Z) + d\eta(Z, X)\eta(Y) \\ &\quad + \frac{1}{3}dF(X, Y, AZ) - \frac{5}{3}dF(AX, AY, AZ) - \frac{1}{3}dF(AX, Y, Z) - \frac{1}{3}dF(X, AY, Z). \end{aligned}$$

Compare [\(3.22\)](#) with [\(3.28\)](#) to derive

$$(3.29) \quad 3dF(AX, AY, AZ) + dF(X, Y, AZ) + dF(AX, Y, Z) + dF(X, AY, Z) = 2(d\eta \wedge \eta)(X, Y, Z)$$

Using [\(2.26\)](#) and [\(3.25\)](#), we derive from [\(3.29\)](#) the following

$$(3.30) \quad dF(AX, AY, AZ) + dF(X, Y, AZ) = \eta(X)d\eta(Y, Z) + \eta(Y)d\eta(Z, X).$$

Substitute [\(3.29\)](#) and [\(3.30\)](#) into [\(3.28\)](#) to get

$$(3.31) \quad N(X, Y, Z) = -\frac{4}{3}dF(AX, AY, AZ) + (d\eta \wedge \eta)(X, Y, Z),$$

which, in particular, shows that the Nijenhuis tensor N is totally skew symmetric.

Applying [\(3.30\)](#) to [\(3.14\)](#) we obtain

$$\begin{aligned} (3.32) \quad g((\nabla_X^g A)Y, Z) &= \frac{1}{3}dF(X, Y, Z) - \frac{1}{6}\eta(Y)d\eta(Z, AX) + \frac{1}{3}\eta(X)d\eta(Y, AZ) + \frac{1}{6}\eta(Z)d\eta(Y, AX) \\ &= -\frac{1}{3}dF(AX, AY, Z) + \frac{1}{6}\eta(Z)d\eta(Y, AX) - \frac{1}{2}\eta(Y)d\eta(AZ, X). \end{aligned}$$

We derive from [\(3.32\)](#) that $g((\nabla_X^g A)Y, Z) + g((\nabla_Y^g A)X, Z) = -\frac{1}{2}\eta(Y)d\eta(AZ, X) - \frac{1}{2}\eta(X)d\eta(AZ, Y)$, i.e. the structure is nearly-cosymplectic, $(\nabla_X^g A)X = 0$ if and only if the 1-form η is closed, $d\eta = 0$.

It seems reasonable to consider the following

Definition 3.6. An almost contact metric manifold $(M^{2n+1}, A, g, F, \eta, \xi)$ is said to be almost-nearly cosymplectic if the Levi-Civita covariant derivative of the fundamental 2-form satisfies the following condition

$$(3.33) \quad g((\nabla_X^g A)Y, Z) = -\frac{1}{3}dF(AX, AY, Z) + \frac{1}{6}\eta(Z)d\eta(Y, AX) - \frac{1}{2}\eta(Y)d\eta(AZ, X).$$

In particular, the vector field ξ is Killing and [\(3.31\)](#) holds.

Theorem [3.1](#) yields

Theorem 3.7. Let (M, A, g, F, η, ξ) be an almost contact metric manifold with a fundamental 2-form F considered as a generalized Riemannian manifold (M, G) with a generalized Riemannian metric $G = g + F$. Then (M, G) satisfies the Einstein metricity condition [\(1.1\)](#) with a totally skew-symmetric torsion T if and only if it is an almost-nearly cosymplectic, i.e. [\(3.33\)](#) holds.

The skew-symmetric torsion is determined by the condition

$$\begin{aligned} T(X, Y, Z) &= -\frac{1}{3}dF(X, Y, Z) \\ &= -\frac{1}{4}N(AX, AY, AZ) + \frac{1}{3}\left[\eta(X)d\eta(Y, AZ) + \eta(Y)d\eta(Z, AX) + \eta(Z)d\eta(X, AY)\right]. \end{aligned}$$

The connection is unique given by the formula

$$g(\nabla_X Y, Z) = g(\nabla_X^g Y, Z) - \frac{1}{6}dF(X, Y, Z) + \frac{1}{6}\left[\eta(X)d\eta(Y, Z) + \eta(Y)d\eta(X, Z)\right].$$

The Einstein metricity condition has the form $(\nabla_X G)(Y, Z) = \frac{1}{3} [dF(X, Y, Z) - dF(X, Y, AZ)]$.
 The covariant derivative of g and F are $(\nabla_X g)(Y, Z) = \frac{1}{6} [\eta(Y)d\eta(Z, X) + \eta(Z)d\eta(Y, X)]$,

$$(\nabla_X F)(Y, Z) = \frac{1}{3} [dF(X, Y, Z) - dF(X, Y, AZ)] - \frac{1}{6} [\eta(Y)d\eta(Z, X) + \eta(Z)d\eta(Y, X)].$$

3.6. Almost-nearly cosymplectic structures. We list some elementary properties of the almost-nearly cosymplectic structures defined by (3.33).

First, we have due to (3.26)

Corollary 3.8. *On an almost-nearly cosymplectic manifold the vector field ξ is Killing.*

Corollary 3.9. *If an almost-nearly cosymplectic structure is normal then it is cosymplectic (coKähler), i.e. $d\eta = dF = 0$.*

Proof. An almost contact structure is normal if the almost contact Nijenhuis tensor vanishes, $N^{ac} = 0$ which, in view of (2.29), gives $N(X, Y, Z) = -\eta(Z)d\eta(X, Y)$. Now, (3.31) yields $d\eta = 0$ and $dF(AX, AY, AZ) = 0$. The last equality together with (3.25) implies $dF = 0$. $\square$

Remark 3.10. *An almost-nearly cosymplectic structure is never contact because of (3.21). Indeed if $d\eta = F$ then $d\eta$ is of type (1,1) with respect to A which contradicts (3.21).*

Suppose an almost-nearly cosymplectic structure has closed 1-form η , $d\eta = 0$. Then (3.21) yields that η is ∇^g -parallel, $\nabla^g \eta = 0$. Then the distribution $H = \text{Ker}(\eta)$ is involutive and therefore it is Frobenius integrable. The integral submanifold N^{2n} is a nearly Kähler manifold due to (3.33). Hence, an almost-nearly cosymplectic manifold with a closed 1-form η is locally a product of a nearly Kähler manifold with the real line.

More general, we have $\mathcal{L}_\xi \eta = d\eta(\xi, \cdot) = 0$ due to (3.21). Since ξ is a Killing vector field, we calculate using (3.26), (3.25) and (3.32) that

$$(3.34) \quad (L_\xi F)(X, Y) = (\nabla_\xi^g F)(X, Y) - g(\nabla_X^g \xi, AY) + g(\nabla_Y^g \xi, AX) = d\eta(Y, AX).$$

Suppose M^{2n+1} is compact and the almost contact metric structure is regular, i.e. the $2n$ -dimensional quotient space M^{2n+1}/ξ is a smooth manifold. If in addition $d\eta = 0$ then (3.34) and (3.32) imply that M^{2n+1} is a trivial circle bundle over the Nearly Kähler manifold M^{2n+1}/ξ . Conversely, start with a Nearly Kähler manifold N^{2n} we obtain on $M^{2n+1} \times \mathbb{R}^+$ an almost-nearly cosymplectic structure taking $\eta = dt$.

3.7. Almost paracontact metric structures. Almost paracontact metric structure is defined in Section 4.6. Equation (3.14) together with equation (2.34) implies the equalities

$$(3.35) \quad \begin{aligned} (\nabla_X^g \eta)Y &= g(\nabla_X^g \xi, Y) = -g((\nabla_X^g A)AY, \xi) = -\frac{1}{3}dF(X, AY, \xi) + \frac{1}{6}dF(AX, Y, \xi); \\ d\eta(X, Y) &= -\frac{1}{6}dF(AX, Y, \xi) - \frac{1}{6}dF(X, AY, \xi), \\ d\eta(X, \xi) &= 0, \quad d\eta(AX, Y) = d\eta(X, AY). \end{aligned}$$

With the help of (3.35) we write the condition (3.10) in the form

$$(3.36) \quad \begin{aligned} N(X, Y, Z) &= \frac{1}{3}dF(X, Y, AZ) + \frac{1}{3}dF(AX, Y, Z) + \frac{1}{3}dF(X, AY, Z) + \frac{1}{3}dF(AX, AY, AZ) \\ &\quad - \eta(X)d\eta(Y, Z) - \eta(Y)d\eta(Z, X) + \eta(Z)d\eta(X, Y) \end{aligned}$$

Now Theorem 3.1 leads to the following

acnikap

Theorem 3.11. *Let (M, A, g, F, η, ξ) be an almost paracontact metric manifold with a fundamental 2-form F considered as a generalized Riemannian manifold $(M, G = g + F)$. Then (M, G) satisfies the Einstein metricity condition ^{metein} (1.1) with a totally skew-symmetric torsion T if and only if the condition ^{ff2} (3.14) holds*

The skew-symmetric torsion is determined by ^{tordfnew} (3.11), the connection is unique given by the formula ^{newnb1} (3.13). The covariant derivatives of g and F are determined by ^{skew0} (3.12).

Proof. We show that ^{ff2} (3.14) implies ^{skew1AP} (3.36) and then apply Theorem ^{pp1} 3.1.

The general formula from ^{Zam} [43, Proposition 2.4], in our notation, reads

ZamAP

$$(3.37) \quad 2g((\nabla_X^g A)Y, Z) = dF(X, Y, Z) + dF(X, AY, AZ) - N^{apc}(Y, Z, AX) \\ + [\eta(AY, Z) + d\eta(Y, AZ)]\eta(X) + d\eta(AY, X)\eta(Z) - d\eta(AZ, X)\eta(Y).$$

Substitute ^{ff2} (3.14) into ^{ZamAP} (3.37) and notice $N^{apc}(Y, Z, AX) = N(Y, Z, AX)$, we get

ZamkAP

$$(3.38) \quad N(Y, Z, AX) = \frac{1}{3}dF(X, Y, Z) + \frac{1}{3}dF(X, AY, AZ) + \frac{1}{3}dF(AX, Y, AZ) + \frac{1}{3}dF(AX, AY, Z) \\ + 2d\eta(AY, Z)\eta(X) + d\eta(AY, X)\eta(Z) - d\eta(AZ, X)\eta(Y).$$

The defining equation ^{hu1} (2.16) of the Nijenhuis tensor together with the properties listed in ^{apcon} (2.34) and ^{ki1AP} (3.35) imply $N(X, Y, \xi) = -d\eta(AX, AY) = -d\eta(X, Y)$. Now, we easily get the equivalences of ^{ZamkAP} (3.38) and ^{skew1AP} (3.36). An application of Theorem ^{pp1} 3.1 completes the proof. $\square$

In view of Theorem ^{acnikap} 3.11 it seems reasonable to give the following

Definition 3.12. *An almost paracontact metric manifold (M, g, η, ξ, A) is called an almost paracontact NGT manifold with torsion if the following condition holds*

$$g((\nabla_X^g A)Y, Z) = \frac{1}{3}dF(X, Y, Z) + \frac{1}{3}dF(X, AY, AZ) - \frac{1}{6}dF(AX, Y, AZ) - \frac{1}{6}dF(AX, AY, Z).$$

On an almost paracontact NGT manifold the equalities ^{ki1AP} (3.35) hold, the Nijenhuis tensor is given by ^{skew1AP} (3.36) and the Nijenhuis tensor $N(AX, AY, AZ)$ is totally skew-symmetric determined by

$$N(AX, AY, AZ) = \frac{1}{3}dF(AX, AY, Z) + \frac{1}{3}dF(X, AY, AZ) + \frac{1}{3}dF(AX, Y, AZ) + \frac{1}{3}dF(X, Y, Z) \\ + 2\eta(Z)d\eta(AX, Y) + 2\eta(Y)d\eta(AZ, X) + 2\eta(X)d\eta(AY, Z).$$

An example can be obtained as follows, start with a Nearly paraKähler manifold N^{2n} one obtains on $M^{2n+1} \times \mathbb{R}^+$ an almost paracontact NGT structure taking $\eta = dt$.

Acknowledgments. S.Ivanov is visiting University of Pennsylvania, Philadelphia. S.I. thanks UPenn for providing the support and an excellent research environment during the final stage of the paper. S.I. is partially supported by Contract DFNI I02/4/12.12.2014 and Contract 168/2014 with the Sofia University "St.Kl.Ohridski". M.Z. was partially supported by and the project EUROWEB, and by Serbian Ministry of Education, Science, and Technological Development, Projects 174012...

REFERENCES

Agr

[1] I. Agricola, *The Srni lectures on non-integrable geometries with torsion*, Arch. Math. (Brno) 42 (2006), suppl., 584. 7

BBDG

[2] K. Becker, M. Becker, K. Dasgupta, P.S. Green, *Compactifications of heterotic theory on non-Kähler complex manifolds: I*, JHEP 0304 (2003), 007. 3

BBE

[3] K. Becker, M. Becker, K. Dasgupta, P.S. Green, E. Sharpe, *Compactifications of heterotic strings on non-Kähler complex manifolds: II*, Nuclear Phys. B 678 (2004), 19–100. 3

- BBDP** [4] K. Becker, M. Becker, K. Dasgupta, S. Prokushkin, *Properties of heterotic vacua from superpotentials*, Nuclear Phys. **B 666** (2003), 144–174. [3](#)
- y4** [5] K. Becker, M. Becker, J.-X. Fu, L.-S. Tseng, S.-T. Yau, *Anomaly cancellation and smooth non-Kähler solutions in heterotic string theory*, Nuclear Phys. **B 751** (2006), 108–128. [3](#), [7](#)
- BSethi** [6] K. Becker, S. Sethi, *Torsional heterotic geometries*, Nuclear Phys. **B 820** (2009), 1–31. [3](#), [7](#)
- Blair** [7] D.E. Blair, *Riemannian geometry of contact and symplectic manifolds*. Second edition. Progress in Mathematics, 203. Birkhäuser Boston, Inc., Boston, MA, 2010. [9](#), [14](#)
- But** [8] J.-B. Butruille, *Twistors and 3-symmetric spaces*, Proc. London Math. Soc. **96** (2008), 738–766. [3](#), [13](#)
- Ein1** [9] A. Einstein, *Die Grundlagen der allgemeinen Relativitätstheorie*, Ann. der Physik, 49, 769, 1916. [2](#)
- Ein** [10] A. Einstein, *The Meaning of Relativity*, Princeton Univ. Press, 1953. [2](#), [10](#)
- Eis** [11] L.P. Eisenhart, *Generalized Riemannian spaces I*, Nat. Acad. Sci. USA, 37, (1951), 311–315. [8](#)
- FIMU** [12] M. Fernandez, S. Ivanov, V. Munos, L. Ugarte, *Nearly hypo structures and compact nearly Kähler 6-manifolds with conical singularities*, J. London Math. Soc. (2) **78** (2008) 580–604. [3](#), [13](#)
- FosH** [13] Lorenzo Foscolo, Mark Haskins, *New G2 holonomy cones and exotic nearly Kähler structures on the 6-sphere and the product of a pair of 3-spheres*, arXiv:1501.07838. [3](#)
- FI** [14] Th. Friedrich, S. Ivanov, *Parallel spinors and connections with skew-symmetric torsion in string theory*, Asian J. Math. **6** (2002), 3003–336. [3](#), [7](#), [8](#), [10](#), [13](#)
- Fried** [15] A. Friedmann, *Über die Krmmung des Raumes*, Zeitschrift für Physik 10 (1922), 377–386. [2](#)
- Gau** [16] P. Gauduchon, *Hermitian connections and Dirac operators*, Boll. Un. Mat. Ital. B (7) **11**(2), Suppl., 257–288 (1997). [8](#)
- GKMW** [17] J.P. Gauntlett, N. Kim, D. Martelli, D. Waldram, *Fivebranes wrapped on SLAG three-cycles and related geometry*, JHEP **0111** (2001), 018. [3](#), [7](#)
- GMPW** [18] J.P. Gauntlett, D. Martelli, S. Pakis, D. Waldram, *G-structures and wrapped NS5-branes*, Commun. Math. Phys. **247** (2004), 421–445. [3](#)
- GMW** [19] J.P. Gauntlett, D. Martelli, D. Waldram, *Superstrings with intrinsic torsion*, Phys. Rev. **D 69** (2004), 086002, 27 pp. [3](#), [7](#)
- GKP** [20] S.B. Giddings, S. Kachru, J. Polchinski, *Hierarchies from fluxes in string compactifications*, Phys. Rev. **D 66** (2002), 106006, 16 pp. [3](#)
- GPap** [21] J. Gillard, G. Papadopoulos, D. Tsimpis, *Anomaly, fluxes and (2,0) heterotic-string compactifications*, JHEP **0306** (2003), 035. [3](#), [7](#)
- Gr1** [22] A. Gray, *Nearly Kähler manifolds*, J. Differential Geom. **4** (1970) 283–309. [3](#), [13](#)
- Gr2** [23] A. Gray, *Riemannian manifolds with geodesic symmetries of order 3*, J. Differential Geom. **7** (1972) 343–369. [3](#), [13](#)
- Gr3** [24] A. Gray, *The structure of nearly Kähler manifolds*, Math. Ann. **223** (1976) 233–248. [3](#), [13](#)
- GLNP** [25] K.-P. Gemmer, O. Lechtenfeld, C. Nölle, A.D. Popov, *Yang-Mills instantons on cones and sine-cones over nearly Kähler manifolds*, JHEP 09 (2011) 103. [3](#), [13](#)
- Ham** [26] R. T. Hammond, *The necessity of torsion in gravity*, Gen Relativ Gravit (2010) **42**:23452348. [3](#)
- Hlav** [27] V. Hlavaty, *Geometry of Einsteins Unified Field Theory*, P. Noordhoff Ltd, Groningen, Holland, 1957. [11](#)
- Iv** [28] S. Ivanov, *Connection with torsion, parallel spinors and geometry of Spin(7) manifolds*, Math. Res. Lett., **11** (2004), 171–186. [7](#)
- IZ** [29] S. Ivanov, S. Zamkovoy, *Para-Hermitian and Para-Quaternionic manifolds*, Diff. Geom. Appl. **23** (2005), 205–234. [8](#), [9](#), [13](#), [14](#)
- Ki** [30] V. Kirichenko, *K-spaces of maximal rank*, Mat. Zam. **22** (1977) 465–476 (Russian). [3](#), [13](#)
- LNP** [31] O. Lechtenfeld, C. Nölle, A.D. Popov, *Heterotic compactifications on nearly Kähler manifolds*, JHEP 09 (2010) 074. [3](#), [13](#)
- Lema** [32] G. Lemaître, *L’univers en expansion*, Ann. Soc. Sci. Brux. 47A, 49 (1927). [2](#)
- KN** [33] S. Kobayashi, K. Nomizu, *Foundations of Differential Geometry*. volI 1963, vol. II 1967 [7](#)
- LMof** [34] J. Legare, J. W. Moffat, *Geodesic and Path Motion in the Nonsymmetric Gravitational Theory*, Gen.Rel.Grav.28:1221-1249,1996 . [2](#), [10](#)
- Mof** [35] J.W. Moffat, *Regularity Theorems in the Nonsymmetric Gravitational Theory*, arXiv:gr-qc/9504009. [2](#), [10](#), [11](#)
- MNS** [36] A. Moroianu, P.A. Nagy, U. Semmelmann, *Unit Killing vector fields on nearly Kähler manifolds*, Internat. J. Math. **16** (2005), no. 3, 281–301. [3](#), [13](#)
- N1** [37] P.-A. Nagy, *Nearly Kähler geometry and Riemannian foliations*, Asian J. Math. 6 (2002), no. 3, 481504. (Reviewer: Maria Falcitelli) 53C15 (53C12 53C28) [3](#), [13](#)
- N2** [38] P.-A. Nagy, *On nearly-Kähler geometry*, Ann. Global Anal. Geom. **22** (2002), no. 2, 167–178. [3](#), [13](#)

- | | |
|------|--|
| Po1 | [39] A.D. Popov, <i>Hermitian-Yang-Mills equations and pseudo-holomorphic bundles on nearly Kähler and nearly Calabi-Yau twistor 6-manifolds</i> , Nucl.Phys. B 828 (2010), 594-624. 3 , 13 |
| PS | [40] A.D. Popov and R.J. Szabo, <i>Double quiver gauge theory and nearly Kähler flux compactifications</i> , JHEP 02 (2012) 033. 3 , 13 |
| Str | [41] A. Strominger, <i>Superstrings with torsion</i> , Nuclear Phys. B 274 (1986), 253–284. 3 , 7 , 8 |
| Yano | [42] K. Yano, <i>Differential geometry on complex and almost complex spaces</i> , Pergamon Press, New York, 1965. 3 , 13 |
| Zam | [43] S. Zamkovoy, <i>Canonical connections on paracontact manifolds</i> , Ann. Global Anal. Geom. 36 (2009), no. 1, 37-60. 3 , 10 , 17 |

(Ivanov) UNIVERSITY OF SOFIA "ST. KL. OHRIDSKI", FACULTY OF MATHEMATICS AND INFORMATICS, BLVD. JAMES BOURCHIER 5, 1164 SOFIA, BULGARIA; INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES

AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, DRL 209 SOUTH 33RD STREET PHILADELPHIA, PA 19104-6395

E-mail address: ivanovsp@fmi.uni-sofia.bg

(Zlatanović) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND MATHEMATICS, UNIVERSITY OF NIŠ, VIŠEGRADSKA 33, 18000 NIŠ, SERBIA

E-mail address: zlatmilan@yahoo.com